

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtbOriginal article <http://dx.doi.org/10.1016/j.apjtb.2016.11.023>Chemical composition of essential oil of *Thymus vulgaris* collected from Saudi Arabian marketAbdulrahman Khazim Al-Asmari^{1*}, Md Tanwir Athar¹, Ahmed Abdullah Al-Faraidy², Mohammed Salim Almuhaiza³¹Department of Research Center, Prince Sultan Military Medical City, Riyadh, Saudi Arabia²Department of Psychiatry, Prince Sultan Military Medical City, Riyadh, Saudi Arabia³Department of Dentistry, Prince Sattam Bin Abdulaziz University, Al-Kharj, Saudi Arabia

ARTICLE INFO

Article history:

Received 2 Mar 2016

Received in revised form 28 Mar, 2nd

revised form 14 Apr 2016

Accepted 20 Jun 2016

Available online 25 Nov 2016

Keywords:

Essential oil

GC–MS

Saudi Arabian herbs

Thymus vulgaris

ABSTRACT

Objective: To focus on the analysis of chemical constituents of the *Thymus vulgaris* L. (locally known as “Zaitra” or “Za'atar”; Family: Lamiaceae) which is available in the market of Saudi Arabia.**Methods:** The Zaitra oil was analyzed by gas chromatography-mass spectrometry. The mass spectra were compared with the standard spectra available in National Institute Standard and Technique library.**Results:** The results indicated that the *Thymus* oil is composed of many chemical compounds including α pinene, thymol and caryophyllene which are biologically active and also used in various diseases.**Conclusions:** It can be concluded that the *Thymus vulgaris* due to presence of many bioactive compounds can be used as a new potential source of medicine for the treatment of various types of illness.

1. Introduction

In the modern era of pharmaceutical science, the extent of the degree of interest has been shifted more towards the herbal products as compared to the classical or synthetic products due to better affordability, acceptability and compatibility with the human physiology and minimal side effects [1]. The interests in herbal products for the medicines, perfumery, cosmetics and food additives are increasing among common people, patients and physicians as evident from an increased market of herbal medicine [2]. Among all these herbal drugs, the essential/volatile oils have received substantial attraction due to their significant biological usefulness [3].

The composition of the aromatic and volatile oils are very much influenced by genotype (species, cultivar, clone, ecotype), ecological (geographical origin, climate condition, soil composition) and technological (cultivation, types of collection, storage of crude material and processing technique) factors [4]. For these reasons, plant of the same species, but from the different contexts can express different features and chemical compositions. In this study attempt was made to characterize the chemicals present in the locally marketed *Thymus vulgaris* L. (Lamiaceae) (*T. vulgaris*).

T. vulgaris is an annual plant with grassy appearance that grows in many parts of the world. It is commonly used in folk medicine for its expectorant, antitussive, antibroncholytic, anti-spasmodic, antihelminthic, carminative and diuretic properties. It is also used as culinary herbs and is a well-known source of flavoring agent [5].

The plant contains aromatic smell and warm pungent taste. The fragrance of this plant is due to essential oil, which gives flavoring value for culinary purposes as well as its medicinal properties [6]. The studies showed that *T. vulgaris* has carminative, antimicrobial, muscle relaxant [6] and antioxidant properties [7,8].

*Corresponding author: Abdulrahman Khazim Al-Asmari, Director, Department of Research Centre, Prince Sultan Military Medical City, Post Box 7897 (775s), Riyadh, Saudi Arabia.

Tel: +966 1 14777714, ext. 25100

Fax: +966 1 4786601

E-mail: abdulrahman.alasmari@gmail.com

Foundation Project: Supported by Prince Sultan Military Medical City, Riyadh, Saudi Arabia (Grant Number # APX-2014-034).

Peer review under responsibility of Hainan Medical University. The journal implements double-blind peer review practiced by specially invited international editorial board members.

Keeping in view the past published work [3,4,7,9–12], on the medically important aspect of this plant, we evaluated the chemical composition of *T. vulgaris*. Additionally, the rationale behind undertaking this project was to compare what are the similarities and differences in bioactive components of the plant marketed in Saudi Arabia and rest of the world due to climate/geographical changes. Since one of the major problems associated with the herbal products and dietary supplements is its quality control [13], the data obtained by this studies may be used as a tool to assess the quality of *Thymus* leaves.

It's worth noting that the *T. vulgaris* which we have studied are not necessarily grown in the Kingdom of Saudi Arabia. It may be imported from other geographical locations of the world. To the best of our knowledge, no one has characterized the chemical composition of *T. vulgaris* marketed and used by the vast majority of the Saudi population. This study is the first of its kind to demonstrate the presence of bioactive components in *T. vulgaris* that is available in Saudi Arabia.

2. Materials and methods

2.1. Plant material

The dried leaves of *T. vulgaris* were collected from the local market in Riyadh, Saudi Arabia in December 2015. It was identified by Dr. Abuzer Ali, Department of Natural Products and Alternative Medicine, Damman University, and the voucher specimen (voucher number P/M/014) has been deposited to Herbal Unit, Research Center, Prince Sultan Military Medical City, Riyadh, Saudi Arabia. The leaves were coarsely powdered and used for analysis.

2.2. Essential oil extraction by steam distillation

A Clevenger apparatus was used to extract volatile oil. Thirty grams of coarsely powdered leaves were taken in a 1000 mL flask, and 750 mL of water was added. The flask was heated on heating mantle. The volatile oil was extracted and it was evaporated along with the water vapor, which were passed through condenser and oil was accumulated. Distillation was stopped when the difference in successive readings of the oil volume remained constant. The oil was drained followed by passing over anhydrous sodium sulfate. The oil was filtered through 0.22 $\mu\text{mol/L}$ filter paper, and kept at 4 °C in sealed vials in dark.

2.3. Instrumentation

Gas chromatography-mass spectrometry (GC–MS) analysis of the *Thymus* oil was carried out on a GC system (Agilent 7890A series, USA) as reported earlier [14]. The instrument was equipped with split/splitless injector. The auto-sampler was attached to an apolar HP-5MS (5% phenyl polymethyl siloxane) capillary column (30 m \times 0.25 mm, i.d. 0.25 μm film thickness) and fitted to Mass Detector (Agilent 5975C series, USA). Helium was used as carrier gas with flow rate of 1 mL/min. Split ratio was maintained at 1:20, injector temperature at 250 °C, detector temperature at 300 °C. The column temperature was kept at 60 °C for 2 min followed by linear programming from 60 to 250 °C (at 2 °C/min), and kept isothermal for 2 min. The

transfer line was heated at 280 °C. Mass spectra were acquired in scan mode (70 eV) in range 50–550 m/z . Two microliter of 1000 $\mu\text{L/L}$ oil sample diluted in hexane was injected. The components of the oil were separated and the chromatogram obtained was identified by comparing the mass spectra to those from National Institute of Standards and Technology (NIST) libraries.

2.4. Identification of chemical constituents

The spectra obtained from the GC–MS were compared with the database of NIST which contains more than 62000 standard spectra. The spectra of the compound obtained from the *Thymus* oil was compared with the spectrum of the known compounds which is stored in the NIST library.

3. Results

3.1. GC–MS optimization

GC–MS parameters were optimized through adjusting the column, temperature program, and split ratio. The best result obtained from HP-5MS column that showed higher resolution and less analysis time. Finally, the optimized GC–MS analysis condition was developed for specific analysis of *T. vulgaris* as stated in instrumentation section.

3.2. Compound identification

The GC–MS fingerprint profile of *T. vulgaris* was obtained and showed in Figure 1. By analyzing the results, hydro-distilled oil of *T. vulgaris* showed the presence of 24 chemical compounds. The compounds, its retention time, and area percentage are shown in Table 1.




Figure 1. GC–MS chromatogram of *T. vulgaris* analyzed on GC–MS (Agilent, USA) using a capillary column (5-MS) attached with mass detector. The chromatogram showed the presence of chemical components found in the *T. vulgaris*.

Table 1

List of the compounds present in the *Thymus* oil analyzed by GC–MS to show the retention time and the area percentage of each compound.

No.	Compound	Retention time (min)	Percentage (%)
1	Furan, tetra hydro-3-methyl	2.317	12.19
2	Cyclohexane	2.546	0.13
3	α -Pinene	6.546	0.18
4	Camphene	6.998	0.13
5	β -Myrcene	8.491	0.15
6	3-Octanol	8.663	0.22
7	Carene	9.389	0.27
8	<i>p</i> -Cymene	9.704	2.77
9	<i>o</i> -Cymene	9.841	0.39
10	γ -Terpinene	11.060	1.18
11	1,6-Octadien-3-ol, 3,7-dimethyl-	12.828	0.38
12	Endo-Borneol	15.563	0.90
13	Terpinen-4-ol	16.090	0.31
14	α -Terpineol	16.811	0.28
15	Thymol	21.354	3.82
16	<i>o</i> -Thymol	22.018	38.71
17	Phenol, 2-methyl-5-(1-methylethyl) acetate	24.610	0.34
18	Caryophyllene	26.349	0.91
19	Humulene	31.385	0.22
20	2-Aminopyrimidine-1-oxide	32.277	0.14
21	(-)-Spathulenol	32.609	0.13
22	Caryophyllene oxide	32.781	0.51
23	Isoaromadendrene epoxide	35.613	0.11
24	Cyclo propane carboxamide	40.013	0.11

A number of compounds found in essential oil of *T. vulgaris* were reported to have strong biological activities. The bioactive compounds including the molecular formula, molecular weight, chemical structure, and reported biological activity are presented in Table 2.

Table 2

Details of bioactive compounds found in the *Thymus* oil after GC–MS analysis along with formula, molecular weight and reported activity of each bioactive compound.

No.	Compound	Molecular formula	Molecular weight	Reported activity	Reference
1	α -Pinene	C ₁₀ H ₁₆	136.24	Antioxidant, antiinflammatory, chondroprotective	[15,16]
2	Camphene	C ₁₀ H ₁₆	136.24	Hypolipidemic, hepatoprotective	[17,18]
3	β -Myrcene	C ₁₀ H ₁₆	136.24	Antioxidant, antiulcer, anticancer	[19–21]
4	Carene	C ₁₀ H ₁₆	136.24	Antioxidant, antiinflammatory	[22–24]
5	<i>p</i> -Cymene	C ₁₀ H ₁₄	134.21	Antioxidant, antimicrobial	[25,26]
6	γ -Terpinene	C ₁₀ H ₁₆	136.24	Antimicrobial, antioxidant	[27]
7	α -Terpineol	C ₁₀ H ₁₈ O	154.25	Antimicrobial	[28]
8	Thymol	C ₁₀ H ₁₄ O	150.22	Antimicrobial, antioxidant	[29]
9	Caryo-phyllene	C ₁₅ H ₂₄	204.36	Local anesthesia, antimicrobial	[30]
10	Humulene	C ₁₅ H ₂₄	204.36	Larvicidal, anticancer	[31,32]

4. Discussion

The quality control of the herbal drug is a big challenge in all around the world. World Health Organization has issued guidelines to validate the herbs that are used for medicinal and therapeutics purposes. Saudi Arabia is exporting a number of herbs which are used as food products as well as for the therapeutic purposes. There is a need to develop the system through which the quality of the herbs can be evaluated in terms of presence or absence of chemical constituent.

GC is an ideal technique for the separation and identification of the herbs containing the volatile and semi volatile constituents. MS attached with the GC has an added advantage through which each separated constituent can be fragmented and the pattern of fragmentation can be compared with the spectra available in the database. This gives the information of the available constituent present in the herbs without using the chemical/biological marker of the plants. Finally, all the available constituents can be separated, fragmented and characterized simultaneously.

The GC–MS of the *Thymus* oil resulted in the identification of 24 compounds. Chemical compounds found in *T. vulgaris* were terpenes and aromatic compounds. The major compounds were thymol, camphene, caryophyllene, humulene, α -terpeniol and para-cymene. Volatile compounds investigated in this study were found to be consistent with those of previously published studies in which volatile compounds were identified by different methods [3,4,7,9].

The high percentage of the thymol found in the Saudi market leaves can be seen as the desirable concentration of the phenol that could have value in the traditional market.

The analyses of this work reveal that the hydro-distilled oil of *T. vulgaris* has a number of components. These bioactive components, present in Zaitra act as a drug for various diseases. The essential oil of *T. vulgaris* reveals the presence of 24 bioactive components from GC–MS technique, and each component has specific functions and acts as a drug for various diseases. Thus from this study it can be concluded that the *T. vulgaris* may serve as a new potential source of medicines due to the presence of these phytochemicals and bioactive compounds. Further, the functional studies of these bioactive compounds will further be examined using *in vivo* models.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

The research was supported by the Prince Sultan Military Medical City, Riyadh, Saudi Arabia (Grant Number # APX-2014-034).

References

- [1] Al-Asmari AK, Al-Elaiwi AM, Athar MT, Tariq M, Al Eid A, Al-Asmari SM. A review of hepatoprotective plants used in Saudi traditional medicine. *Evid Based Complement Alternat Med* 2014; 2014: 890842.
- [2] Musthaba M, Baboota S, Athar MT, Thajudeen KY, Ahmed S, Ali J. Patented herbal formulations and their therapeutic applications. *Recent Pat Drug Deliv Formul* 2010; 4: 231-44.

- [3] Miladi H, Slama RB, Mili D, Zouari S, Bakhrouf A, Ammar E. Essential oil of *Thymus vulgaris* L. and *Rosmarinus officinalis* L.: gas chromatography–mass spectrometry analysis, cytotoxicity and antioxidant properties and antibacterial activities against foodborne pathogens. *Nat Sci* 2013; **5**(6): 729-39.
- [4] Mancini E, Senatore F, Del Monte D, De Martino L, Grulova D, Scognamiglio M, et al. Studies on chemical composition, antimicrobial and antioxidant activities of five *Thymus vulgaris* L. essential oils. *Molecules* 2015; **20**: 12016-28.
- [5] Javanmardi J, Khalighi A, Kashi A, Bais H, Vivanco J. Chemical characterization of basil (*Ocimum basilicum* L.) found in local accessions and used in traditional medicines in Iran. *J Agric Food Chem* 2002; **50**: 5878-83.
- [6] Boskabady MH, Aslani MR, Kiani S. Relaxant effect of *Thymus vulgaris* on guinea-pig tracheal chains and its possible mechanism(s). *Phytother Res* 2006; **20**: 28-33.
- [7] Lee SJ, Umamo K, Shibamoto T, Lee KG. Identification of volatile components in basil (*Ocimum basilicum* L.) and thyme leaves (*Thymus vulgaris* L.) and their antioxidant properties. *Food Chem* 2005; **91**: 131-7.
- [8] Schwarz K, Ernst H, Ternes W. Evaluation of antioxidative constituents from thyme. *J Sci Food Agric* 1996; **70**: 217-23.
- [9] Golmakani MT, Rezaei K. Comparison of microwave-assisted hydrodistillation with the traditional hydrodistillation method in the extraction of essential oils from *Thymus vulgaris* L. *Food Chem* 2008; **109**: 925-30.
- [10] Rota MC, Herrera A, Martínez RM, Sotomayor JA, Jordán MJ. Antimicrobial activity and chemical composition of *Thymus vulgaris*, *Thymus zygis* and *Thymus hyemalis* essential oils. *Food Control* 2008; **19**: 681-7.
- [11] Hashmi LS, Hossain MA, Weli AM, Al-Riyami Q, Al-Sabahi JN. Gas chromatography–mass spectrometry analysis of different organic crude extracts from the local medicinal plant of *Thymus vulgaris* L. *Asian Pac J Trop Biomed* 2013; **3**: 69-73.
- [12] Mardafkan N, Iranmanesh M, Larijani K, Mahasti P, Nazari F, Zojaji M. Chemical components and antibacterial activities of essential oils obtained from Iranian local *Lavandula officinalis* and *Thymus vulgaris* against pathogenic bacteria isolated from human. *J Food Biosci Technol* 2015; **5**: 31-6.
- [13] Musthaba SM, Athar MT, Baboota S, Ahmed S. HPTLC analysis of neem oil extract in herbal dosage form. *Asian J Chem* 2011; **23**: 385-7.
- [14] Al-Asmari AK, Albalawi SM, Athar MT, Khan AQ, Al-Shahrani H, Islam M. *Moringa oleifera* as an anti-cancer agent against breast and colorectal cancer cell lines. *PLoS One* 2015; **10**: e0135814.
- [15] Kovač J, Šimunović K, Wu Z, Klančnik A, Bucar F, Zhang Q, et al. Antibiotic resistance modulation and modes of action of (–)- α -pinene in *Campylobacter jejuni*. *PLoS One* 2015; **10**: e0122871.
- [16] Rufino AT, Ribeiro M, Judas F, Salgueiro L, Lopes MC, Cavaleiro C, et al. Anti-inflammatory and chondroprotective activity of (+)- α -pinene: structural and enantiomeric selectivity. *J Nat Prod* 2014; **77**: 264-9.
- [17] Vallianou I, Hadzopoulou-Cladaras M. Camphene, a plant derived monoterpene, exerts its hypolipidemic action by affecting SREBP-1 and MTP expression. *PLoS One* 2016; **11**(1): e0147117.
- [18] Kim S, Choi Y, Choi S, Choi Y, Park T. Dietary camphene attenuates hepatic steatosis and insulin resistance in mice. *Obesity* 2014; **22**: 408-17.
- [19] Ciftci O, Ozdemir I, Tanyildizi S, Yildiz S, Oguzturk H. Antioxidative effects of curcumin, β -myrcene and 1,8-cineole against 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced oxidative stress in rats liver. *Toxicol Ind Health* 2011; **27**: 447-53.
- [20] Bonamin F, Moraes TM, Dos Santos RC, Kushima H, Faria FM, Silva MA, et al. The effect of a minor constituent of essential oil from *Citrus aurantium*: the role of β -myrcene in preventing peptic ulcer disease. *Chem Biol Interact* 2014; **212**: 11-9.
- [21] Lee JH, Lee K, Lee DH, Shin SY, Yong Y, Lee YH. Anti-invasive effect of β -myrcene, a component of the essential oil from *Pinus koraiensis* cones, in metastatic MDA-MB-231 human breast cancer cells. *J Korean Soc Appl Biol Chem* 2015; **58**: 563-9.
- [22] Aazza S, Lyoussi B, Miguel MG. Antioxidant and anti-acetylcholinesterase activities of some commercial essential oils and their major compounds. *Molecules* 2011; **16**: 7672-90.
- [23] Ocete MA, Risco S, Zarzuelo A, Jimenez J. Pharmacological activity of the essential oil of *Bupleurum gibraltaricum*: anti-inflammatory activity and effects on isolated rat uteri. *J Ethnopharmacol* 1989; **25**: 305-13.
- [24] Romeo FV, De Luca S, Piscopo A, Poiana M. Antimicrobial effect of some essential oils. *J Essent Oil Res* 2008; **20**: 373-9.
- [25] de Oliveira TM, de Carvalho RBF, da Costa IHF, de Oliveira GAL, de Souza AA, de Lima SG, et al. Evaluation of *p*-cymene, a natural antioxidant. *Pharm Biol* 2015; **53**: 423-8.
- [26] Silva M, Ribeiro FP, Medeiros MA, Sampaio PA, Silva Y, Silva MT, et al. The vasorelaxant effect of *p*-cymene in rat aorta involves potassium channels. *ScientificWorldJournal* 2015; **2015**: 458080.
- [27] Masoum S, Mehran M, Ghaheri S. Identification of potential antioxidant compounds in the essential oil of thyme by gas chromatography with mass spectrometry and multivariate calibration techniques. *J Sep Sci* 2015; **38**: 410-7.
- [28] Park SN, Lim YK, Freire MO, Cho E, Jin D, Kook JK. Antimicrobial effect of linalool and α -terpineol against periodontopathic and cariogenic bacteria. *Anaerobe* 2012; **18**: 369-72.
- [29] Kavooosi G, Dadfar SM, Purfard AM. Mechanical, physical, antioxidant, and antimicrobial properties of gelatin films incorporated with thymol for potential use as nano wound dressing. *J Food Sci* 2013; **78**: E244-50.
- [30] Dahham SS, Tabana YM, Iqbal MA, Ahamed MB, Ezzat MO, Majid AS, et al. The anticancer, antioxidant and antimicrobial properties of the sesquiterpene β -caryophyllene from the essential oil of *Aquilaria crassna*. *Molecules* 2015; **20**: 11808-29.
- [31] Govindarajan M, Benelli G. α -Humulene and β -elemene from *Syzygium zeylanicum* (Myrtaceae) essential oil: highly effective and eco-friendly larvicides against *Anopheles subpictus*, *Aedes albopictus*, and *Culex tritaeniorhynchus* (Diptera: Culicidae). *Parasitol Res* 2016; **115**(7): 2771-8.
- [32] Hanušová V, Skálová L, Ambrož M, Králová V, Langhansová L, Matoušková P. The effect of *Myrica rubra* essential oil and its components α -humulene and trans-nerolidol on adhesion and apoptosis of colorectal cancer cells. *Cancer Cell Microenviron* 2015; **2**: e1058.