



HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtbReview article <http://dx.doi.org/10.1016/j.apjtb.2016.11.028>

Anticonvulsant effects of medicinal plants with emphasis on mechanisms of action



Zahra Rabiei*

Medical Plants Research Center, Shahrekord University of Medical Science, Shahrekord, Iran

ARTICLE INFO

Article history:

Received 10 Aug 2016

Received in revised form 19 Sep, 2nd

revised form 19 Oct 2016

Accepted 22 Nov 2016

Available online 18 Dec 2016

Keywords:

Epileptic seizure

Anticonvulsant

Medicinal plants

Oxidative stress

ABSTRACT

Epilepsy is a disorder in brain in which clusters of nerve cells, or neurons, occasionally signal abnormally and cause strange emotions, sensations, and behavior, or sometimes muscle spasms, convulsions, and loss of consciousness. Neurotransmitters in central nervous system greatly affect and play a very important part in neuronal excitability. Traditional treatments are still a component of health care system in many communities despite the fact that well-established alternatives are available. In this review article, we addressed epilepsy and its treatments with emphasis on medicinal plants and introduction of antiepileptic plants and their action mechanisms. Relevant articles published since 2010 were retrieved using the search terms including epileptic seizure, anticonvulsant, medicinal plants, and oxidative stress. Most plants/herbal preparations that are ethnomedicinally used to treat epilepsy or those which have been tested for anticonvulsant activity were reported. Overall, the results of the published articles show that the symptoms of epilepsy seizure can be inhibited or treated by active ingredients derived from medicinal plants.

1. Introduction

Epileptic seizure syndromes can be due to a wide variety of causes, including genetic, developmental, or acquired ones. Seizures mainly occur suddenly without warning, have short duration (a few seconds or minutes), and stop by themselves [1]. Epileptic seizures are considered to be the most common neurologic symptoms in different human populations and remain the most common neurological condition involving people at any age. At any time, fifty million worldwide are estimated to have a diagnosis of epilepsy [2].

Epileptic seizures are seizure events that occur due to excessive, abnormally synchronized, localized, or widely distributed neuronal electrical discharges [3].

An epileptic seizure is an episode of neurologic dysfunction due to abnormal neuronal firing obviously occurring clinically via changes in sensory perception, motor control, behavior, or autonomic function [4].

2. Types of epileptic seizures

Seizures have two main types, *i.e.* focal or partial seizures and generalized seizures. In focal seizures, only one part of the brain, occasionally called the 'focus' of the seizures, is affected. Focal seizure may affect a large part of one hemisphere or only a small area of a lobe but generalized seizures occur when seizure activity is widespread in the brain's left and right hemispheres and the affected people become unconscious (except in myoclonic seizures), though for a few seconds [5].

3. Basic mechanisms of epilepsy

Seizure initiation is characterized by two parallel conditions: 1) high-frequency bursts of action potentials, and 2) hyper synchronization of a neuronal population [6].

The bursting activity due to the neuronal membrane's relatively prolonged depolarization occurs because of influx of extracellular Ca^{++} , which results in influx of Na^+ , opening of voltage-dependent Na^+ channels, and generation of repetitive action potentials. The hyperpolarizing potential is mediated by Cl^- influx and gamma-aminobutyric acid (GABA) receptors, or by K^+ efflux, according to the cell type [7]. GABA is a type of the brain's inhibitory neurotransmitter, which effectively prevents the brain from sending messages [7].

*Corresponding author: Zahra Rabiei, Medical Plants Research Center, Shahrekord University of Medical Science, Shahrekord, Iran.

Tel: +98 9132815431

E-mail: zahrarabiei@ymail.com

Foundation Project: Funded by the Research and Technology Deputy of the Shahrekord University of Medical Sciences (grant number: 2672).

Peer review under responsibility of Hainan Medical University. The journal implements double-blind peer review practiced by specially invited international editorial board members.

GABA interneurons may result in paradoxical facilitation of certain types of epileptic discharges in some models. Drugs that cause increase in synaptic GABA by inhibition of GABA catabolism or reuptake, are considered effective anticonvulsants, including benzodiazepines, which improve GABA binding to the GABA receptor, leading to increased frequency of chloride channel openings [8]. Some GABA synthesis inhibitors are able to cause seizures, including thiosemicarbazide, 4-deoxyypyridoxine, isoniazid, and L-allylglycine [8].

GABA, the major inhibitory neurotransmitter, interacts with two key subtypes of receptors: GABA_A and GABA_B. GABA_A receptors are found postsynaptically, but GABA_B receptors are found presynaptically and can therefore modulate synaptic release. GABA_A receptors are permeable to Cl⁻ ions in the brain in adulthood; Cl⁻ influx, upon activation, hyperpolarizes the membrane and inhibits action potentials. Therefore, GABA_A receptor agonists, including barbiturates and benzodiazepines, suppress seizure activity. GABA_B receptors are related to second messenger systems but not Cl⁻ channels, and result in attenuation of transmitter release because of their presynaptic location [8].

Glutamate is a type of amino acid and a major excitatory neurotransmitter in the brain. Glutamate released from synapses is taken up by astrocytes under normal conditions, and is rapidly converted to the non-excitotoxic amino acid glutamine by glutamine synthetase [9].

The ionotropic *N*-methyl-D-aspartate (NMDA), α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid/kainate, and metabotropic glutamate receptor-mediated mechanisms are involved in epileptic seizures [9].

Excitatory glutamatergic mechanisms are involved in acute, transient, evoked seizures as well as long-term, adaptive cellular plasticity related to epileptogenesis in chronic epilepsy models. Glutamate exerts its excitatory effects through ligand-gated ion channels (NMDA and non-NMDA receptors) in order to increase sodium and calcium conductance [10].

Neuronal (EAAC1) and glial glutamate transporters facilitate reuptake of glutamate and aspartate after synaptic release. Down-regulation of glutamate transporters can be compatible with enhanced excitatory activity [11].

4. Epilepsy and oxidative stress

Oxidative stress leads to cellular damage and functional cellular disruption and can subsequently cause cell death through oxidation of biomolecules such as lipids, proteins, and nucleotides [12].

Seizure generation can be associated with the homeostatic imbalance between antioxidants and oxidants. Oxidative stress has been described as an imbalance between generation and elimination of reactive oxygen species (ROS) and reactive nitrogen species [13].

ROS levels are relatively well regulated to do significant functions including autophagy, cell division, chemical signaling, and mitogen-activated protein kinase signaling and apoptosis. Because of the highly reactive nature of this molecule, the ROS is tightly regulated. ROS-induced mitochondrial dysfunction is frequently seen following seizures throughout epileptogenesis [14].

Epileptic seizure causes initiation of remarkable influx of calcium through voltage-gated and NMDA-dependent ion channels that escalate intracellular ions and bring about biochemical cascades. High levels of intracellular calcium can

induce ROS [15]. ROS may be scavenged by certain enzymatic antioxidant defense systems such as superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase, and peroxiredoxins and non-enzymatic ones such as vitamin C, vitamin E, and reduced glutathione (GSH) [12]. Antioxidant therapies to reduce oxidative stress have attracted much attention in treatment for epilepsy.

5. Epilepsy and inflammation

Experimental findings in rodents have indicated that seizures cause inflammatory mediators to increase in brain regions involved in generating epileptic activity [16]. Direct anti-inflammatory treatments have been found to suppress some types of epileptic seizures. Inflammatory processes may occur prior to the onset of epilepsy in humans, potentially contributing etiopathogenetically to incidence of spontaneous seizures. A rapid-onset inflammatory response is triggered in glia by seizures that have been induced by chemoconvulsants or electrical stimulation [17].

The over-expressing cytokines, *e.g.* interleukin-6 (IL-6) and tumor necrosis factor- α , inside astrocytes have been reported to cause age-dependent neurological dysfunctions such as decreased seizure threshold and spontaneous seizure frequency [18].

Inflammatory cytokines such as IL-1 β and high-mobility group box 1, activate IL-1 receptor type I and Toll-like receptor 4, respectively. IL-1 receptor/Toll-like receptor signaling can regulate neuronal excitability, including inhibition of Ca²⁺ channels outward current, alteration of synaptic transmission, and decrease in GABA production [19].

6. Antiepileptic drugs

Most epileptic seizures are controlled by drug therapy, especially anticonvulsants. Treatments for seizures are based on anticonvulsant medication, although there are various choices of anticonvulsant drugs with different seizure types and epileptic syndromes [20]. Patients with newly diagnosed epilepsy who need treatment can start treatment with standard anticonvulsants including carbamazepine, valproic acid/valproate semisodium, phenytoin, phenobarbital, or more recently with gabapentin, oxcarbazepine, lamotrigine, or topiramate [21].

The type of prescribed treatment depends on different factors including the severity and frequency of the seizures as well as age, overall health, and medical history. Accurate diagnosis of epilepsy type is essential to choose the best treatment [21].

Conventional antiepileptic drugs may block sodium channels or improve GABA function. Different antiepileptic drugs have multiple or uncertain mechanisms of action [22]. Next to the voltage-gated sodium channels and components of the GABA system, their targets include GABA_A receptors, the GABA transporter 1, and GABA transaminase. Other targets include voltage-gated calcium channels, SV2A, and $\alpha_2\delta$ [23].

Through blocking sodium or calcium channels, antiepileptic drugs decrease the release of excitatory glutamate, which increases in epilepsy, and also GABA [23]. This is possibly a side effect or the actual action mechanism of some of the antiepileptic drugs, because GABA can directly or indirectly act proconvulsively [24].

Most of the drugs currently being used have unpleasant side effects and unpredictable pharmacological actions; therefore, it

is necessary to search for newer drugs with fewer or no side effects and predictable pharmacological action because treatment of epilepsy is a long-term process and drug taking is discontinued gradually for about six months [25].

Nature is a rich source of biological and chemical varieties. The peerless and complicated structures of natural products cannot be figured out easily by chemical synthesis [26]. Medicinal plants play an important role in treating neurological disease such as

Alzheimer's disease [27–31], brain ischemia, reperfusion [32–35], and other degenerative diseases [36–38].

In reality, current interest in traditional medicine has resulted in quick development and investigation of many remedies applied in various ethnic groups across the world. The data on the type of extract, essential oil and active components, mechanisms of action, methods and reference related to the plants which have been tested or reported for anticonvulsant properties are summarized in Table 1.

Table 1

Antiepileptic medicinal plants or compounds of plant origin.

Compounds	Methods (induced seizures)	Mechanism of action	Reference
<i>Withania somnifera</i> methanolic extract	Unilateral hippocampal injection of kainite (1 mg/kg) in male rats	Ameliorated spatial memory deficit in Y-maze	[39]
<i>Trichosanthes dioica</i> Roxb fruits aqueous extract	Delivering electroshock (50 mA) for 0.2 s through a pair of ear clip electrodes, PTZ (80 mg/kg) <i>i.p.</i> injection induces tonic-clonic convulsions in mice	Activity against generalized tonic-clonic and cortical focal seizures	[40]
<i>Ficus platyphylla</i> methanol extract	PTZ (37.5 mg/kg) <i>i.p.</i> for a total of 13 convulsant injections in mice, learning performance was tested in a two-way shuttle-box	Affinity for undifferentiated glutamate receptors, affinity for the 3H-GABA binding assay, decrease the K ⁺ -stimulated glutamate release from rat hippocampal slices	[40]
<i>Feretia apodanthera</i> Del lyophilized aqueous extract	Administered PTZ (30 mg/kg; <i>i.p.</i>) 22nd injection in mice, behavioral tests include elevated plus-maze and T-maze tests	Decreased brain MDA levels, increased brain GSH levels, increase of AChE and BChE activity in brain	[41]
<i>Nigella sativa</i> oil	Electroshock and PTZ in mice	Potent antioxidant actions	[42]
<i>Psidium guajava</i> (guava) leaves ethanolic extract	Delivering electroshock (50 mA) for 0.2 s through a pair of ear clip electrodes and PTZ (70 mg/kg) <i>i.p.</i> injection induces tonic-clonic convulsions in mice	Selectively inhibit NMDA receptor	[43]
<i>Trachyspermum ammi</i> (L.) methanol extract	Strychnine (4 mg/kg) <i>i.p.</i> injection-induced seizure in rats	Excite GABA responses mainly by stimulating human GABA _A receptors and increasing the chloride ion channel opening	[44]
<i>Zingiber officinale</i> (ginger) rhizomes hydroethanolic extract	Administrated PTZ into the tail vein	Antioxidant activity, inhibit NO production, reduce inducible nitric oxide synthase	[45]
<i>Anacyclus pyrethrum</i> root hydroalcoholic extract	Administrated <i>i.p.</i> injection of PTZ (60 mg/kg), behavioral tests include elevated plus-maze and passive avoidance tests	Increase in GSH levels of brain, decreased MDA levels of brain, increased AChE and BChE activity in brain	[46]
<i>Valeriana officinalis</i> root aqueous extract	Amygdala-kindled rats	Existence of adenosine ligand(s) in the valerian aqueous extract and activation of A1 adenosine system, binding to GABA receptors	[47]
<i>Panax ginseng</i> root butanol extract	Seizures were induced by administration of 40 mg/kg pilocarpine hydrochloride, magnetic resonance imaging study of the rat brain	Increased hippocampal volume, increase in T ₂ relaxation time in rat hippocampus, entorhinal cortex, piriform cortex, amygdala, thalamus	[48]
<i>Pimpinella anisum</i> essential oil	Administrated PTZ (120 mg/kg) in rat	Inhibited production of dark neurons, inhibited induction of long-term potentiation in hippocampal slices	[49]
<i>Cyperus rotundus</i> Linn. essential oil	MES induced convulsion in rats	Inhibit voltage-dependant Na ⁺ channels, block glutamatergic excitation mediated by the NMDA receptor	[50]
<i>Zhumeria majdae</i> essential oil and methanolic extract	PTZ (110 mg/kg) and MES models in mice	Inhibit voltage-dependent Na ⁺ channels, block glutamatergic excitation mediated by the NMDA receptor	[51]
<i>Angelica archangelica</i> Linn. roots essential oil	PTZ (80 mg/kg) and MES models in mice	Block glutamatergic excitation	[52]
<i>Cymbopogon winterianus</i> Jowitt essential oils	Pilocarpine-induced convulsions (350 mg/kg <i>i.p.</i>) administrated PTZ in mice	GABAergic mechanisms, deteriorated autoregulation of glutamate release	[53]
<i>Rosmarinus officinalis</i> essential oils	Administrated PTZ (80 mg/kg) in mice	Elevate GABA levels in the midbrain region	[54]
<i>Ocimum basilicum</i> essential oils	Administrated PTZ (80 mg/kg) in mice	Modulate glutamate activation expression, reduction of the ACh-evoked release, direct interaction with the NMDA receptor complex	[54]
<i>Mentha spicata</i> essential oils	Administrated PTZ (80 mg/kg) in mice	GABAergic mechanisms	[54]

Table 1 (continued)

Compounds	Methods (induced seizures)	Mechanism of action	Reference
<i>Lavandula angustifolia</i> essential oils	Administrated PTZ (80 mg/kg) in mice	Modulate glutamate activation expression, reduction of the ACh-evoked release, direct interaction with the NMDA receptor complex	[54]
<i>Mentha piperita</i> essential oils	Administrated PTZ (80 mg/kg) in mice	GABAergic mechanisms	[54]
<i>Origanum dictamnus</i> essential oils	Administrated PTZ (80 mg/kg) in mice	GABAergic mechanisms	[54]
<i>Origanum vulgare</i> essential oils	Administrated PTZ (80 mg/kg) in mice	GABAergic mechanisms	[54]
<i>Mentha pulegium</i> essential oils	Administrated PTZ (80 mg/kg) in mice	GABAergic mechanisms	[54]
Turmeric methanolic extract	40 mmol/L PTZ in zebrafish larvae, PTZ (50, 100 mg/kg) infusion in tail vein in mice	Higher lipophilicity and easily cross the blood–brain barrier, suppress oxidative DNA damage and lipid peroxidation	[55]
α -Asarone	MES seizure test in mice, pilocarpine administered <i>i.p.</i> (320 and 350 mg/kg), pentylenetetrazole administered (85 mg/kg)	Inhibited pyrogallol auto-oxidation, α -asarone displays 5 units of superoxide dismutase-like activity, inhibit the hydroperoxide dependent oxidation of glutathione	[56]
<i>Acorus calamus</i> Linn aqueous extract	PTZ (80 mg/kg) and MES models in mice	Block NMDA receptors	[57]
<i>Bunium persicum</i> essential oil and methanolic extract	PTZ (110 mg/kg) induced convulsion in NMRI male mice, MES models	GABAergic mechanisms	[58]
Phytol (a constituent of chlorophyll)	Pilocarpine hydrochloride (400 mg/kg <i>i.p.</i>) in mice	Muscarinic activation and alterations in AChE activity in hippocampus	[59]
Soy extract	Administrated PTZ (40 mg/kg <i>i.p.</i>) for 14 days or a single injection of a high dose of PTZ (90 mg/kg in ovariectomized rat)	Phytoestrogens of soy affect seizure severity	[60]
<i>Zizyphus jujuba</i> hydroalcoholic extract	Administrated PTZ (60 mg/kg <i>i.p.</i>), MES in rat	Increase in brain GSH, decreased brain MDA levels, increased brain AChE and BChE activity	[61]
<i>Emblica officinalis</i> hydroalcoholic extract	Administrated PTZ (60 mg/kg) in rat	Decreased brain MDA levels, increase in brain GSH levels	[62]
Naringin (bioflavonoid present in the grapefruit)	Administrated PTZ (60 mg/kg) in rat	Increased brain GSH levels, decreased levels of MDA, attenuated the high brain levels of tumor necrosis factor- α	[63]
<i>Anisomeles malabarica</i> (flavonoids fraction from the leaves)	MES, administrated PTZ (50 mg/kg) in rat	Decreased tonic hindlimb extension phase and extensor/flexion ratio in MES model	[64]
Curcumol (from <i>Rhizoma Curcumae</i>)	Hippocampal neurons in culture	Facilitation of GABA _A Rs by curcumol in hippocampal neurons, facilitation of recombinant GABA _A Rs, enhancement of phasic GABAergic inhibition by curcumol in hippocampal slices, enhancement of tonic GABAergic inhibition by curcumol in hippocampal slices	[65]
Hydroalcoholic extract of citrus flower	Administrated PTZ (90 mg/kg) in rat	Increase in minimal clonic seizure	[66]
Vitexin (a flavonoid)	Vitexin administered intracerebroventricularly, administered PTZ (90 mg/kg <i>i.p.</i>)	Vitexin is a ligand for benzodiazepine receptors, exerts anticonvulsant effects through a GABA _A benzodiazepine receptor	[67]
Quercetin (a flavonoid)	Administration of PTZ every other day (35 mg/kg <i>i.p.</i> , 15 injections in total)	Decreased brain MDA levels, increased sulfhydryl in brain	[68]
<i>Centella asiatica</i> (gotu kola)	Injection of pentylenetetrazol (60 mg/kg <i>i.p.</i>)	Increased AChE activity, elevated levels of ACh	[69]
Curcumin	Administration of PTZ (35 mg/kg) in mice	Increased brain norepinephrine level, reduced total nitrite levels of brain, reduced AChE activity	[70]
<i>Nigella sativa</i> oil	Injection of a single dose of pilocarpine (380 mg/kg <i>i.p.</i>)	Attenuate the increased NO levels resulting from pilocarpine, attenuate the decrease in hippocampal Na ⁺ , K ⁺ ATPase activity, increase the AChE enzyme	[71]

PTZ: Pentylenetetrazole; MES: Maximal electroshock; MDA: Malondialdehyde; AChE: Acetylcholinesterase; BChE: Butyrylcholinesterase; ACh: Acetylcholine.

7. Discussion

Epilepsy is the second leading neurological disorder after stroke, involving at least 50 million worldwide [72]. Cognitive impairment, dose-related neurotoxicity, and a spectrum of systemic side effects are the main side effects due to antiepileptic drugs [20].

The PTZ kindling model is a commonly used screening model to test anticonvulsive compounds. It exerts action mainly through the *t*-butyl-bicyclo-phosphorothionate/picrotoxin site of the GABA_A receptor. PTZ is a blocker of choice for the GABA_A receptor chloride ionophore complex [73]. It has convulsant effects after repeated or single administration and affects several neurotransmitter systems, such as adenosinergic, GABAergic,

and glutamatergic systems [74]. After PTZ-induced seizures, significant decreases in GSH, cysteine, glutathione disulfide, and protein thiols as well as increases in the protein disulfides and protein carbonyl levels were observed in the mouse cerebral cortex [75].

The MES seizure test, in which tonic hindlimb seizures are induced by bilateral corneal or transauricular electrical stimulation, is believed to predict effectiveness of anticonvulsant drug against generalized tonic-clonic seizures [76]. Local or systemic administration with pilocarpine and kainate in rodents has led to a pattern of repetitive limbic seizures and status epilepticus, lasting for several hours [77].

The drugs used to treat epilepsy may cause certain side effects. The occurrence of side effects depends on the dose of the drug taken, duration of treatment, and type of medication. The side effects are more likely to occur due to taking higher doses of the drugs but tend to mitigate over time because of the body's adjustment to the medication [78]. Natural products and their derivatives comprise over 50% of all the drugs used in clinical settings worldwide [79].

Medical plants can be applied because of their structural diversity and wide spectrum of pharmacological effects in contrast to common synthetic antiepileptic drugs. Available evidence suggests that many herbal medicines may cause adverse effects in people suffering from seizure [79].

It can be concluded that studies on active compounds in the plant-based extracts may be important to identify chemical compounds for development of antiepileptic drugs in the future.

Conflict of interest statement

I declare that I have no conflict of interest.

Acknowledgments

This research was funded by the Research and Technology Deputy of the Shahrekord University of Medical Sciences (grant number: 2672).

References

- [1] Yalçın AD, Kaymaz A, Forta H. Childhood occipital epilepsy: seizure manifestations and electroencephalographic features. *Brain Dev* 1997; **19**(6): 408-13.
- [2] Fisher RS, van Emde Boas W, Blume W, Elger C, Genton P, Lee P, et al. Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* 2005; **46**(4): 470-2.
- [3] Nakagawa E, Aimi Y, Yasuhara O, Tooyama I, Shimada M, McGeer PL, et al. Enhancement of progenitor cell division in the dentate gyrus triggered by initial limbic seizures in rat models of epilepsy. *Epilepsia* 2000; **41**(1): 10-8.
- [4] Scott BW, Wojtowicz JM, Burnham WM. Neurogenesis in the dentate gyrus of the rat following electroconvulsive shock seizures. *Exp Neurol* 2000; **165**(2): 231-6.
- [5] Lopes da Silva FH, Blanes W, Kalitzin SN, Parra J, Suffczynski P, Velis DN. Dynamical diseases of brain systems: different routes to epileptic seizures. *IEEE Trans Biomed Eng* 2003; **50**(5): 540-8.
- [6] Richerson GB, Wu Y. Role of the GABA transporter in epilepsy. *Adv Exp Med Biol* 2004; **548**: 76-91.
- [7] Schuler V, Löscher C, Blanchet C, Klix N, Sansig G, Klebs K, et al. Epilepsy, hyperalgesia, impaired memory, and loss of pre- and postsynaptic GABA_B responses in mice lacking GABA_B(1). *Neuron* 2001; **31**(1): 47-58.
- [8] Gale K. GABA and epilepsy: basic concepts from preclinical research. *Epilepsia* 1991; **33**: S3-12.
- [9] During MJ, Spencer DD. Extracellular hippocampal glutamate and spontaneous seizure in the conscious human brain. *Lancet* 1993; **341**(8861): 1607-10.
- [10] Doherty J, Dingledine R. The roles of metabotropic glutamate receptors in seizures and epilepsy. *Curr Drug Targets CNS Neurol Disord* 2002; **1**(3): 251-60.
- [11] Vespa P, Prins M, Ronne-Engstrom E, Caron M, Shalmon E, Hovda DA, et al. Increase in extracellular glutamate caused by reduced cerebral perfusion pressure and seizures after human traumatic brain injury: a microdialysis study. *J Neurosurg* 1998; **89**(6): 971-82.
- [12] Sudha K, Rao AV, Rao A. Oxidative stress and antioxidants in epilepsy. *Clin Chim Acta* 2001; **303**: 19-24.
- [13] Gluck MR, Jayatilake E, Shaw S, Rowan AJ, Haroutunian V. CNS oxidative stress associated with the kainic acid rodent model of experimental epilepsy. *Epilepsy Res* 2000; **39**(1): 63-71.
- [14] Frantseva MV, Perez Velazquez JL, Tsoraklidis G, Mendonca AJ, Adamchik Y, Mills LR, et al. Oxidative stress is involved in seizure-induced neurodegeneration in the kindling model of epilepsy. *Neuroscience* 2000; **97**(3): 431-5.
- [15] Coyle JT, Puttfarcken P. Oxidative stress, glutamate, and neurodegenerative disorders. *Science* 1993; **262**(5134): 689-95.
- [16] Marchi N, Granata T, Janigro D. Inflammatory pathways of seizure disorders. *Trends Neurosci* 2014; **37**(2): 55-65.
- [17] Kwon YS, Pineda E, Auvin S, Shin D, Mazarati A, Sankar R. Neuroprotective and antiepileptogenic effects of combination of anti-inflammatory drugs in the immature brain. *J Neuroinflammation* 2013; **10**: 30.
- [18] Virta M, Hurme M, Helminen M. Increased plasma levels of pro- and anti-inflammatory cytokines in patients with febrile seizures. *Epilepsia* 2002; **43**(8): 920-3.
- [19] Kalueff AV, Lehtimäki KA, Ylinen A, Honkaniemi J, Peltola J. Intranasal administration of human IL-6 increases the severity of chemically induced seizures in rats. *Neurosci Lett* 2004; **365**(2): 106-10.
- [20] Perucca E, Gram L, Avanzini G, Dulac O. Antiepileptic drugs as a cause of worsening seizures. *Epilepsia* 1998; **39**(1): 5-17.
- [21] Temkin NR. Antiepileptogenesis and seizure prevention trials with antiepileptic drugs: meta-analysis of controlled trials. *Epilepsia* 2001; **42**(4): 515-24.
- [22] Ketter T, Post RM, Theodore WH. Positive and negative psychiatric effects of antiepileptic drugs in patients with seizure disorders. *Neurology* 1998; **53**(5 Suppl 2): S53-67.
- [23] Rogawski MA, Löscher W. The neurobiology of antiepileptic drugs. *Nat Rev Neurosci* 2004; **5**(7): 553-64.
- [24] Kammerer M, Rassner MP, Freiman TM, Feuerstein TJ. Effects of antiepileptic drugs on GABA release from rat and human neocortical synaptosomes. *Naunyn-Schmiedeberg's Arch Pharmacol* 2011; **384**(1): 47-57.
- [25] Ortinski P, Meador KJ. Cognitive side effects of antiepileptic drugs. *Epilepsy Behav* 2004; **5**: S60-5.
- [26] Samuels N, Finkelstein Y, Singer SR, Oberbaum M. Herbal medicine and epilepsy: proconvulsive effects and interactions with antiepileptic drugs. *Epilepsia* 2008; **49**(3): 373-80.
- [27] Rabiei Z, Rafeiean-Kopaei M, Heidarian E, Saghaei E, Mokhtari S. Effects of *Zizyphus jujube* extract on memory and learning impairment induced by bilateral electric lesions of the nucleus basalis of Meynert in rat. *Neurochem Res* 2014; **39**(2): 353-60.
- [28] Rabiei Z, Rafeiean-Kopaei M, Mokhtari S, Alibabaei Z, Shahrani M. The effect of pretreatment with different doses of *Lavandula officinalis* ethanolic extract on memory, learning and nociception. *Biomed Aging Pathol* 2014; **4**(1): 71-6.
- [29] Rahnema S, Rabiei Z, Alibabaei Z, Mokhtari S, Rafeiean-Kopaei M, Deris F. Anti-amnesic activity of *Citrus aurantium* flowers extract against scopolamine-induced memory impairments in rats. *Neurol Sci* 2015; **36**(4): 553-60.
- [30] Rabiei Z, Hojjati M, Rafeiean-Kopaei M, Alibabaei Z. Effect of *Cyperus rotundus* tubers ethanolic extract on learning and memory in animal model of Alzheimer. *Biomed Aging Pathol* 2013; **3**(4): 185-91.

- [31] Rabiei Z, Mokhtari S, Asgharzade S, Gholami M, Rahnama S, Rafieian-Kopaei M. Inhibitory effect of *Thymus vulgaris* extract on memory impairment induced by scopolamine in rat. *Asian Pac J Trop Biomed* 2015; **5**(10): 845-51.
- [32] Rabiei Z, Rafieian-Kopaei M. Neuroprotective effect of pretreatment with *Lavandula officinalis* ethanolic extract on blood-brain barrier permeability in a rat stroke model. *Asian Pac J Trop Med* 2014; **7**: S421-6.
- [33] Rabiei Z, Bigdeli MR, Rasoulia B, Ghassempour A, Mirzajani F. The neuroprotection effect of pretreatment with olive leaf extract on brain lipidomics in rat stroke model. *Phytomedicine* 2012; **19**(10): 940-6.
- [34] Rabiei Z, Bigdeli MR, Rasoulia B. Neuroprotection of dietary virgin olive oil on brain lipidomics during stroke. *Curr Neurovasc Res* 2013; **10**(3): 231-7.
- [35] Jivad N, Rabiei Z. Review on herbal medicine on brain ischemia and reperfusion. *Asian Pac J Trop Biomed* 2015; **5**(10): 789-95.
- [36] Moradi MT, Rafieian-Kopaei M, Imani-Rastabi R, Nasiri J, Shahrani M, Rabiei Z, et al. Antispasmodic effects of yarrow (*Achillea millefolium* L.) extract in the isolated ileum of rat. *Afr J Tradit Complement Altern Med* 2013; **10**(6): 499-503.
- [37] Rahimian GA, Rabiei Z, Tahmasebi B, Rafieian-Kopaei M, Ganji F, Rahimian R, et al. Comparing the combined effect of garlic and mint extract with metronidazole in *Helicobacter pylori* treatment. *Iran J Pharm Sci* 2013; **9**(3): 63-70.
- [38] Rabiei Z, Rafieian-Kopaei M, Mokhtari S, Shahrani M. Effect of dietary ethanolic extract of *Lavandula officinalis* on serum lipids profile in rats. *Iran J Pharm Res* 2014; **13**(4): 1295-301.
- [39] Narenjkar J, Shoja-Shafiee F, Roghani M. The effect of *Withania somnifera* alcoholic extract on learning and memory disturbance in a model of temporal lobe epilepsy in the rat. *J Basic Clin Pathophysiol* 2014; **3**(1): 13-8.
- [40] Singh P, Garg VK, Sharma PK, Gupta S. Antiepileptic activity of aqueous extract of *Tricosanthes dioica* Roxb. *Asian J Plant Sci Res* 2012; **2**(1): 45-7.
- [41] Taiwe GS, Moto FC, Ayissi ER, Ngoupaye GT, Njapdounke JS, Nkantchoua GC, et al. Effects of a lyophilized aqueous extract of *Feretia apodanthera* Del. (Rubiaceae) on pentylenetetrazole-induced kindling, oxidative stress, and cognitive impairment in mice. *Epilepsy Behav* 2015; **43**: 100-8.
- [42] Jaykare SC, Motghare VM, Padwal SL, Deshmukh VS, Patil JR, Pise HN, et al. Evaluation of anticonvulsant activity of the seed oil extract of *Nigella sativa*: an experimental study. *Hygeia J Drugs Med* 2013; **5**: 21-6.
- [43] Pushpa VH, Shetty KP, Sushma N, Kalabharathi HL, Satish AM. Evaluation of the anticonvulsant activity of ethanol extract of *Psidium guajava* (guava leaves) in albino mice. *Int J Pharm Sci Res* 2014; **5**(10): 4288-92.
- [44] Rajput MA, Khan RA, Feroz Z. Evaluation of antiepileptic activity of the methanol extract of *Trachyspermum ammi* (L.). *Arch Biol Sci* 2013; **65**(3): 815-9.
- [45] Hosseini A, Mirazi N. Acute administration of ginger (*Zingiber officinale* rhizomes) extract on timed intravenous pentylenetetrazol infusion seizure model in mice. *Epilepsy Res* 2014; **108**(3): 411-9.
- [46] Pahuja M, Mehla J, Reeta K, Joshi S, Gupta YK. Root extract of *Anacyclus pyrethrum* ameliorates seizures, seizure-induced oxidative stress and cognitive impairment in experimental animals. *Epilepsy Res* 2012; **98**(2): 157-65.
- [47] Rezvani ME, Roohbakhsh A, Allahtavakoli M, Shamsizadeh A. Anticonvulsant effect of aqueous extract of *Valeriana officinalis* in amygdala-kindled rats: possible involvement of adenosine. *J Ethnopharmacol* 2010; **127**(2): 313-8.
- [48] Suleymanova E, Gulyaev M, Chepurnova N. Ginseng extract attenuates early MRI changes after status epilepticus and decreases subsequent reduction of hippocampal volume in the rat brain. *Epilepsy Res* 2014; **108**(2): 223-31.
- [49] Karimzadeh F, Hosseini M, Mangeng D, Alavi H, Hassanzadeh GR, Bayat M, et al. Anticonvulsant and neuroprotective effects of *Pimpinella anisum* in rat brain. *BMC Complement Altern Med* 2012; **12**: 76.
- [50] Biradar S, Kangralkar V, Mandavkar Y, Thakur M, Chougule N. Antiinflammatory, antiarthritic, analgesic and anticonvulsant activity of *Cyperus* essential oils. *Int J Pharm Pharm Sci* 2010; **2**(4): 112-5.
- [51] Mandegary A, Shariffar F, Abdar M, Arab-Nozari M. Anticonvulsant activity and toxicity of essential oil and methanolic extract of *Zhumeria majdae* Rech, a unique Iranian plant in mice. *Neurochem Res* 2012; **37**(12): 2725-30.
- [52] Pathak S, Wanjari MM, Jain SK, Tripathi M. Evaluation of anti-seizure activity of essential oil from roots of *Angelica archangelica* Linn. in mice. *Indian J Pharm Sci* 2010; **72**(3): 371-5.
- [53] Silva MR, Ximenes RM, da Costa JG, Leal LK, de Lopes AA, Viana GS. Comparative anticonvulsant activities of the essential oils (EOs) from *Cymbopogon winterianus* Jowitt and *Cymbopogon citratus* (DC) Stapf. in mice. *Naunyn Schmiedebergs Arch Pharmacol* 2010; **381**(5): 415-26.
- [54] Koutroumanidou E, Kimbaris A, Kortsaris A, Bezirtzoglou E, Polissiou M, Charalabopoulos K, et al. Increased seizure latency and decreased severity of pentylenetetrazol-induced seizures in mice after essential oil administration. *Epilepsy Res Treat* 2013; **2013**: 532657.
- [55] Orellana-Paucar AM, Serruys AS, Afrikanova T, Maes J, De Borggraeve W, Alen J, et al. Anticonvulsant activity of bisabolene sesquiterpenoids of *Curcuma longa* in zebrafish and mouse seizure models. *Epilepsy Behav* 2012; **24**(1): 14-22.
- [56] Pages N, Maurois P, Delplanque B, Bac P, Stables JP, Tamariz J, et al. Activities of α -asarone in various animal seizure models and in biochemical assays might be essentially accounted for by antioxidant properties. *Neurosci Res* 2010; **68**(4): 337-44.
- [57] Gopalakrishna HN, Pemminati S, Giri S, Shenoy AK, Holla GKS, Nair V, et al. Effect of *Acorus calamus* on electrical and chemical induced seizures in mice. *Int J Appl Biol Pharm Technol* 2010; **1**(2): 465-73.
- [58] Mandegary A, Arab-Nozari M, Ramiar H, Shariffar F. Anticonvulsant activity of the essential oil and methanolic extract of *Bunium persicum* (Boiss). B. Fedtsch. *J Ethnopharmacol* 2012; **140**(2): 447-51.
- [59] Costa JP, Ferreira PB, De Sousa DP, Jordan J, Freitas RM. Anticonvulsant effect of phytol in a pilocarpine model in mice. *Neurosci Lett* 2012; **523**(2): 115-8.
- [60] Ebrahimzadeh Bideskan AR, Hosseini M, Mohammadpour T, Karami R, Khodamoradi M, Nemati Karimoooy H, et al. Effects of soy extract on pentylenetetrazol-induced seizures in ovariectomized rats. *Zhong Xi Yi Jie He Xue Bao* 2011; **9**(6): 611-8.
- [61] Pahuja M, Mehla J, Reeta K, Joshi S, Gupta YK. Hydroalcoholic extract of *Zizyphus jujuba* ameliorates seizures, oxidative stress, and cognitive impairment in experimental models of epilepsy in rats. *Epilepsy Behav* 2011; **21**(4): 356-63.
- [62] Golechha M, Bhatia J, Arya DS. Hydroalcoholic extract of *Embllica officinalis* Gaertn. affords protection against PTZ-induced seizures, oxidative stress and cognitive impairment in rats. *Indian J Exp Biol* 2010; **48**: 474-8.
- [63] Golechha M, Sarangal V, Bhatia J, Chaudhry U, Saluja D, Arya DS. Naringin ameliorates pentylenetetrazol-induced seizures and associated oxidative stress, inflammation, and cognitive impairment in rats: possible mechanisms of neuroprotection. *Epilepsy Behav* 2014; **41**: 98-102.
- [64] Choudhary N, Bijjem KR, Kalia AN. Antiepileptic potential of flavonoids fraction from the leaves of *Anisomeles malabarica*. *J Ethnopharmacol* 2011; **135**(2): 238-42.
- [65] Ding J, Wang JJ, Huang C, Wang L, Deng S, Xu TL, et al. Curcuminol from *Rhizoma Curcumae* suppresses epileptic seizure by facilitation of GABA_A receptors. *Neuropharmacology* 2014; **81**: 244-55.
- [66] Hosseini M, Pkan P, Rakhshandeh H, Aghaie A, Sadeghnia HR, Rahbardar MG. The effect of hydro-alcoholic extract of citrus flower on pentylenetetrazole and maximal electroshock-induced seizures in mice. *World Appl Sci J* 2011; **15**(8): 1104-9.
- [67] Abbasi E, Nassiri-Asl M, Shafeei M, Sheikhi M. Neuroprotective effects of vitexin, a flavonoid, on pentylenetetrazole-induced seizure in rats. *Chem Biol Drug Des* 2012; **80**(2): 274-8.

- [68] Nassiri-Asl M, Moghbelinejad S, Abbasi E, Yonesi F, Haghghi MR, Lotfizadeh M, et al. Effects of quercetin on oxidative stress and memory retrieval in kindled rats. *Epilepsy Behav* 2013; **28**(2): 151-5.
- [69] Visweswari G, Prasad KS, Chetan PS, Lokanatha V, Rajendra W. Evaluation of the anticonvulsant effect of *Centella asiatica* (gotu kola) in pentylenetetrazol-induced seizures with respect to cholinergic neurotransmission. *Epilepsy Behav* 2010; **17**(3): 332-5.
- [70] Choudhary KM, Mishra A, Poroikov VV, Goel RK. Ameliorative effect of curcumin on seizure severity, depression like behavior, learning and memory deficit in post-pentylenetetrazole-kindled mice. *Eur J Pharmacol* 2013; **704**(1): 33-40.
- [71] Ezz HS, Khadrawy YA, Noor NA. The neuroprotective effect of curcumin and *Nigella sativa* oil against oxidative stress in the pilocarpine model of epilepsy: a comparison with valproate. *Neurochem Res* 2011; **36**(11): 2195-204.
- [72] Forsgren L, Beghi E, Oun A, Sillanpää M. The epidemiology of epilepsy in Europe – a systematic review. *Eur J Neurol* 2005; **12**(4): 245-53.
- [73] Velisek L, Kubova H, Pohl M, Stankova L, Mareš P, Schickerova R. Pentylenetetrazol-induced seizures in rats: an ontogenetic study. *Naunyn Schmiedebergs Arch Pharmacol* 1992; **346**(5): 588-91.
- [74] Diehl RG, Smialowski A, Gotwo T. Development and persistence of kindled seizures after repeated injections of pentylenetetrazol in rats and guinea pigs. *Epilepsia* 1984; **25**(4): 506-10.
- [75] Patsoukis N, Zervoudakis G, Panagopoulos NT, Georgiou CD, Angelatou F, Matsokis NA. Thiol redox state (TRS) and oxidative stress in the mouse hippocampus after pentylenetetrazol-induced epileptic seizure. *Neurosci Lett* 2004; **357**(2): 83-6.
- [76] Barton ME, Peters SC, Shannon HE. Comparison of the effect of glutamate receptor modulators in the 6 Hz and maximal electroshock seizure models. *Epilepsy Res* 2003; **56**(1): 17-26.
- [77] Turski WA, Cavalheiro EA, Schwarz M, Czuczwar SJ, Kleinrok Z, Turski L. Limbic seizures produced by pilocarpine in rats: behavioural, electroencephalographic and neuropathological study. *Behav Brain Res* 1983; **9**(3): 315-35.
- [78] Vermeulen J, Aldenkamp AP. Cognitive side-effects of chronic antiepileptic drug treatment: a review of 25 years of research. *Epilepsy Res* 1995; **22**(2): 65-95.
- [79] Gurib-Fakim A. Medicinal plants: traditions of yesterday and drugs of tomorrow. *Mol Aspects Med* 2006; **27**(1): 1-93.